

\*VIRUCIDAL ACTIVITY - This product when used on LEMON DISINFECTANT NEUTRAL CLEANER IS A product per gallon of water. Apply solution with a cloth, Item No. 91101 environmental, inanimate, hard, non-porous surfaces HOSPITAL USE BACTERICIDE AND \*VIRUCIDE AT 4 mob, sponge or hand pump trigger spraver making sure to exhibits effective virucidal activity against HIV-1. HIV-2. OUNCES PER GALLON. MODIFIED IN THE wet all surfaces completely. Let air dry. Prepare a fresh Hepatitis B Virus (HBV), Hepatitis C Virus (HCV), Herpes PRESENCE OF 400 PPM HARD WATER AS CaCO3, solution for each use. Repeat application at weekly Simplex Type 1 (causative agent of fever blisters). Herpes THIS PRODUCT IS A FUNGICIDE AT 4 OUNCES PER intervals or when mildew growth appears. PRECAUTIONARY STATEMENTS

Simplex Type 2 (genital), Influenza A2/Hong Kong, GALLON, MODIFIED IN THE PRESENCE OF 200 PPM Vaccinia, Rotavirus, Human Coronavirus (ATCC VR-740, HARD WATER AS CaCO3. Strain 229E), SARS associated Coronavirus, Bovine Viral Kills Pandemic 2009 H1N1 Influenza A virus (formerly

Diarrhea Virus (BVDV), Pseudorabies, Bovine called swine flu),

Rhinotracheitis, Feline Leukemia, Feline Picornavirus, GENERAL CLEANING - Apply this product to soiled area Avian Influenza A Virus H9N2, Avian Influenza A Virus with a mop, cloth, sponge, hand pump trigger spraver or Causes irreversible eye damage and skin burns. Do not Octyl decyl dimethyl ammonium chloride H3N2, Canine Distemper Virus, Rabies Virus, low pressure coarse spraver. This product can be diluted get in eyes, on skin or on clothing. May be fatal if absorbed Di Paramyxovirus (Mumps) and Porcine Respiratory & at 4 ounces per gallon or applied at full strength. Reproductive Syndrome Virus (PRRSV). Add 4 ounces of Thoroughly wet soiled surface and scrub as necessary. the product per gallon of water. For heavily soiled areas, a **DEODORIZATION** - To deodorize, apply this product as pre-cleaning step is required. Apply solution with a cloth, indicated under the heading DISINFECTION. mop, sponge, hand pump trigger sprayer or low pressure Veterinary Practice/Animal Care/Animal before eating, drinking, or using tobacco.

coarse spraver so as to wet all surfaces thoroughly. Allow Laboratory/Zoos/Pet Shop/Kennels Disinfection excess liquid.

hard non-porous surfaces: equipment not used for animal

STORAGE AND DISPOSAL the surface to remain wet for 10 minutes, then remove Directions: For cleaning and disinfecting the following DO NOT CONTAMINATE WATER. FOOD. OR FEED BY

STORAGE OR DISPOSAL.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS.

DANGER

KILLS HIV-1. HIV-2. HBV. and HCV ON PRE food or water, utensils, instruments, cages, kennels, PESTICIDE STORAGE - Store in a dry place no lower in CLEANED ENVIRONMENTAL SURFACES/OBJECTS stables, catteries, etc. Remove all animals and feed from temperature than 50°F or higher than 120°F. Store in an PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in premises, animal transportation vehicles, crates, etc. area inaccessible to children.

health care settings (Hospitals, Nursing Homes) or other Remove all litter, droppings and manure from floors, walls CONTAINER DISPOSAL - Nonrefillable container. Do of inanimate surfaces/objects with blood or body fluids, Thoroughly clean all surfaces with soap or detergent and available. Triple rinse container promptly after emptying. and in which the surfaces/objects likely to be solied with rinse with water. Saturate surfaces with a use-solution of 4 Triple rinse as follows: Fill container ¼ full with water and continue rinsing eve. Call a poison control center or doctor for treatment microorganisms. blood or body fluids can be associated with the potential ounces of this product or product name per gallon of water recap. Agitate vigorously, Follow Pesticide Disposal advice. or Type 2 (HIV-1 or HIV-2) (associated with AIDS), Hepatitis closed spaces. Do not house animals or employ after the flow begins to drip. Repeat procedure two more immediately with plenty of water for 15-20 minutes. Call poison control center repeated use. equipment until treatment has been absorbed, set or times. Then offer for recycling or reconditioning. If not or doctor for treatment advice, B Virus (HBV), and Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND dried.

brushes, razors, manicure/pedicure tools, clippers, mixture, or rinsate is a violation of Federal Law. If these BLOOD/BODY FLUIDS.

PERSONAL PROTECTION: When handling items soiled scissors, trimmer blades) in a use-solution of 4 ounces of wastes cannot be disposed of by use according to label with blood or body fluids, use disposable latex gloves, this product per gallon of water for at least 10 minutes, instructions, contact your State Pesticide of Rinse instruments thoroughly and dry before reuse. A Environmental Control Agency, or the Hazardous Waste

gowns, masks, and eve coverings, CLEANING PROCEDURES: Blood and other body fluids fresh use-solution must be prepared daily or more often if representative at the nearest EPA Regional Office for

before application of this product.

minutes

other body fluids should be autoclaved and disposed of cause damage to stainless steel or metal instruments. according to local regulations for infectious waste MILDEWSTAT To control mold and mildew (Aspergillus disposal

must be thoroughly cleaned from surfaces and objects the use-solution becomes visibly dirty. NOTE: Plastics guidance. may remain immersed until ready to use. Stainless steel CONTACT TIME: Allow surface to remain wet for 10 shears and instruments must be removed from the usesolution after 10 minutes, rinsed, dried and kept in a clean. DISPOSAL OF INFECTIOUS MATERIALS: Blood and non-contaminated receptacle. Prolonged immersion may

niger) and the odors they cause on pre-cleaned, hard, non-porous inanimate surfaces add 4 ounces of this

available, puncture and dispose in a sanitary landfill.



CLEANER • DISINFECTANT • \* VIRUCIDE • KILLS GERMS • DEODORIZER

0.04.40

## KEEP OUT OF REACH OF CHILDREN. CORROSIVE. ACTIVE INGREDIENTS

	Octyl decyl dimetnyl ammonium chioride
	Dioctyl dimethyl ammonium chloride 0.407%
	Didecyl dimethyl ammonium chloride0.407%
face shield, rubber gloves, and protective clothing.	
Remove contaminated clothing and wash before reuse.	dimethyl benzyl ammonium chloride 1.085%
Wash thoroughly with soap and water after handling and	INERT INGRÉDIENTS:

**KEEP OUT OF REACH OF CHILDREN** DANGER

## FIRST AID

center or doctor. or going for treatment.

If swallowed: Call poison control center or doctor immediately for treatment DECONTAMINATION AGAINST HIV-1, HIV-2, HBV, Disinfection of Barber/Salon Tools Directions: PESTICIDE DISPOSAL - Pesticide wastes are acutely advice. Have person sip a glass of water if able to swallow. Do not induce and HCV OF SURFACES/OBJECTS SOILED WITH Immerse pre-cleaned barber/salon tools (combs, hazardous. Improper disposal of excess pesticide, spray vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

> If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. You may contact 800/535-5053 for emergency medical treatment information. NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. See side panels for additional precautionary statements.

## NET CONTENTS: 1 GALLON (3.785 L)

EPA Reg. No. 1839-168-70925 EPA Est. No. 70925-CA-001 Chemcor Chemical Corporation 13770 Benson Ave., Chino, CA 91710

Lemon Disinfectant Neutral Cleaner is a phosphate-free, BACTERICIDAL STABILITY OF USE-SOLUTION: isolate), Escherichia coli. Escherichia coli (clinical pH neutral formulation designed to provide effective Prepare a fresh use-solution daily or more often if the use- isolate). Fusobacterium necrophorum, Klebsiella cleaning, deodorizing, and disinfection for hospitals, solution becomes visibly dirty, contaminated or diluted, pneumoniae subsp. pneumoniae. Lactobacillus casei nursing homes, households, homes, private residences, Non-Acid Toilet Bowl Disinfection/Cleaner subsp. rhamnosus, Listeria monocytogenes, Pasteurella schools, food processing plants, food service Directions; Remove gross filth prior to disinfection, multocida, Proteus vulgaris, Proteus mirabilis ATCC 9921. establishments, restaurants, transportation terminals, From concentrate; Add 4 ounces per gallon to the toilet Proteus mirabilis ATCC 25933, Salmonella (paratyphi B) office buildings, manufacturing facilities, lodging bowl and mix. Brush thoroughly over exposed surfaces enterica, Salmonella (typhi) enterica, Salmonella establishments, hotels, retail business, and and under the rim, allow to stand for 10 minutes and flush. (typhimurium) enterica, Salmonella (pullorum) enterica, athletic/recreation facilities, where housekeeping is of From use-solution: Empty toilet bowl or urinal and apply Serratia marcescens, Shigella sonnei, Shigella flexneri prime importance in controlling cross-contamination from 4 ounces per gallon use-solution to exposed surfaces, Type 2b, Shigella dysenteriae, Staphylococcus aureus treated surfaces. brush or swab thoroughly under the rim with a cloth, mop, subsp. aureus, Staphylococcus epidermidis, Lemon Disinfectant Neutral Cleaner is a one step neutral sponge or spraver, allow to stand for 10 minutes and flush. Staphylococcus epidermidis (clinical isolate). disinfectant that is effective against a broad spectrum of For heavy duty use: Empty toilet bowl or urinal and Streptococcus pyogenes (Clinical - Flesh Eating Strain bacteria, is virucidal\* (including HIV-1, HIV-2, HBV and apply 16 ounces per gallon use-solution to exposed BIRD M3), Streptococcus pyogenes Group A, HCV) and inhibits the growth of mold and mildew and their surfaces, brush or swab thoroughly under the rim with a Xanthamonas maltophilia (clinical isolate). Vancomycin odors when used as directed. cloth, mop, sponge or sprayer, allow to stand for 10 resistant Enterococcus faecalis (VRE), Methicillin Lemon Disinfectant Neutral Cleaner, when used as minutes and flush. resistant Staphylococcus aureus (MRSA), Vancomycin

a mop, cloth, sponge, hand pump trigger sprayer or low Staphylococcus aureus (CA-MRSA) NRS 384 Genotype the area or carefully protected. After use, all surfaces in Acinetobacter baumannii. Gentamicin resistant countertops, appliances, and tables, add 4 ounces of Acinetobacter baumannii, Levofloxacin resistant

directed, is formulated to disinfect the following hard non- To disinfect food processing premises: floors, walls, intermediate resistant Staphylococcus aureus (VISA), porous inanimate environmental surfaces: floors, walls, and storage areas, add 4 ounces of Lemon Disinfectant Community Associated Methicillin resistant metal surfaces, stainless steel surfaces, glazed porcelain, Neutral Cleaner per gallon of water. For heavily soiled Staphylococcus aureus (CA-MRSA) NRS 123 Genotype glazed ceramic tile, plastic surfaces, vanity tops, shower areas, a pre-cleaning step is required. Apply solution with USA400. Community Associated resistant stalls, bathtubs, and cabinets, Have the product container or label with you when calling a poison control This product deodorizes those areas which generally are pressure coarse sprayer so as to wet all surfaces USA300, Salmonella (entertitidis) enterica, Ampicillin hard to keep fresh smelling, such as garbage storage thoroughly. Allow to remain wet for 10 minutes, then resistant Acinetobacter baumannii, Cefazolin resistant settings in which there is an expected likelihood of soiling and surfaces occupied or traversed by animals. not reuse or refill this container. Offer for recycling, if If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 areas, empty garbage bins and cans, toilet bowls, and remove excess liquid. Before using this product, food Acinetobacter baumannii, Ceftazdime resistant minutes. Remove contact lenses, if present, after the first 5 minutes, then other areas which are prone to odors caused by products and packaging materials must be removed from Acinetobacter baumannii, Ceftriaxone resistant Lemon Disinfectant Neutral Cleaner is formulated to a the area must be thoroughly rinsed with potable water. Acinetobacter baumannii, Tobramycin resistant for transmission of Human Immunodeficiency Virus Type 1 for a period of 10 minutes. Ventilate buildings and other instructions for rinsate disposal. Drain for 10 seconds If on skin or clothing: Take off contaminated clothing. Rinse skin neutral pH and will not dull high-gloss floor finishes with To disinfect restaurant food contact surfaces: Acinetobacter baumannii, Ciprofloxacin resistant This product is an economical concentrate that can be Lemon Disinfectant Neutral Cleaner per gallon of water. Acinetobacter baumannii and Bactrim resistant used with a mop and bucket, trigger spravers or sponge.

For heavily soiled areas, a pre-cleaning step is required. Acinetobacter baumannii. DIRECTIONS FOR USE Apply solution with a cloth, sponge or hand pump trigger FUNGICIDAL ACTIVITY - At the 4 ounce per gallon surface to remain wet for 10 minutes, then remove excess fungi, Trichophyton mentagrophytes (Athlete's Foot on liquid and rinse the surface with potable water. This Fungus) and Candida albicans. Apply solution with a product cannot be used to clean the following food cloth, sponge or hand pump trigger sprayer to hard, nonporous surfaces found in bathrooms, shower stalls, locker

It is a violation of Federal Law to use this product in a sprayer so as to wet all surfaces thoroughly. Allow the dilution, this product is fungicidal against the pathogenic manner inconsistent with its labeling. Lemon Disinfectant Neutral Cleaner is not for critical and semi-critical medical device surfaces. **DISINFECTION** To disinfect inanimate, hard non- contact surfaces; utensils, glassware and dishes,

porous surfaces add 4 ounces of Lemon Disinfectant BACTERICIDAL ACTIVITY - At the 4 ounce per gallon rooms, exercise facilities or other clean, hard non-porous Neutral Cleaner per gallon of water. Apply solution with a dilution. Lemon Disinfectant Neutral Cleaner surfaces commonly contacted by bare feet. Allow the mop, cloth, sponge, hand pump trigger sprayer or low demonstrates effective disinfectant activity against the surface to remain wet for 10 minutes, then remove excess pressure coarse spraver so as to wet all surfaces organisms; Pseudomonas aeruginosa PRD-10, liquid, Diluted product should be applied daily or more thoroughly, Allow to remain wet for 10 minutes, then Salmonella (choleraesuis) enterica. Staphylococcus frequently with heavy facility use. remove excess liquid. For spraver applications, sprav 6-8 aureus, Staphylococcus aureus (clinical isolate). inches from surface, rub with brush, sponge or cloth. Do Bordetella bronchiseptica, Corvnebacterium not breathe spray mist. ammoniagenes. Enterobacter aerogenes. Enterobacter For heavily soiled areas, a pre-cleaning step is required. cloacae. Enterobacter cloacae (clinical isolate) Enterococcus faecalis, Enterococcus faecalis (clinical